

ON LIE IDEALS AND JORDAN LEFT DERIVATIONS
OF PRIME RINGS

MOHAMMAD ASHRAF AND NADEEM-UR-REHMAN

ABSTRACT. Let R be a 2-torsion free prime ring and let U be a Lie ideal of R such that $u^2 \in U$ for all $u \in U$. In the present paper it is shown that if d is an additive mappings of R into itself satisfying $d(u^2) = 2ud(u)$ for all $u \in U$, then $d(uv) = ud(v) + vd(u)$ for all $u, v \in U$.

1. INTRODUCTION

Throughout the present paper R will denote an associative ring with centre Z . Recall that R is prime if $aRb = 0$ implies that $a = 0$ or $b = 0$. As usual $[x, y]$ will denote the commutator $xy - yx$. An additive subgroup U of R is said to be a Lie ideal of R if $[u, r] \in U$ for all $u \in U, r \in R$. An additive mapping $d : R \rightarrow R$ is called a derivation (resp. Jordan derivation) if $d(xy) = d(x)y + xd(y)$, (resp. $d(x^2) = d(x)x + xd(x)$) holds for all $x, y \in R$. Obviously every derivation is a Jordan derivation. The converse need not be true in general. A famous result due to Herstein [8] states that every Jordan derivation on a 2-torsion free prime ring is a derivation. A brief proof of this result is presented in [4]. Further, Awtar [1] generalized this result on Lie ideals.

An additive mapping $d : R \rightarrow R$ is called a left derivation (resp. Jordan left derivation) if $d(xy) = xd(y) + yd(x)$ (resp. $d(x^2) = 2xd(x)$) holds for all $x, y \in R$. Clearly, every left derivation is a Jordan left derivation. Thus, it is natural to question that: Whether every Jordan left derivation on a ring is a left derivation? In the present paper we have shown that the answer to the above question is affirmative in the case when the underlying ring R is 2-torsion free and prime. In fact we have obtained rather a more general result which establish that under appropriate restriction on a Lie ideal U of a 2-torsion free prime ring, every Jordan left derivation on U is a left derivation on U .

2000 *Mathematics Subject Classification*: 16W25, 16N60.

Key words and phrases: Lie ideals, prime rings, Jordan left derivations, left derivations, torsion free rings.

Received August 23, 1999.

2. PRELIMINARY RESULTS

We begin with the following results which will be used extensively to prove our theorem. Lemma 2.1 can be found in [2].

Lemma 2.1. *If $U \not\subseteq Z$ is a Lie ideal of a 2-torsion free prime ring R and $a, b \in R$ such that $aUb = 0$, then $a = 0$ or $b = 0$.*

Lemma 2.2. *Let R be a 2-torsion free ring and let U be a Lie ideal of R such that $u^2 \in U$ for all $u \in U$. If $d : R \rightarrow R$ is an additive mapping satisfying $d(u^2) = 2ud(u)$ for all $u \in U$, then*

- (i) $d(uv + vu) = 2ud(v) + 2vd(u)$, for all $u, v \in U$.
- (ii) $d(uvu) = u^2d(v) + 3uvd(u) - vud(u)$, for all $u, v \in U$.
- (iii) $d(uvw + wvu) = (uw + wu)d(v) + 3uvd(w) + 3wvd(u) - vud(w) - vwd(u)$, for all $u, v, w \in U$.
- (iv) $[u, v]ud(u) = u[u, v]d(u)$, for all $u, v \in U$.
- (v) $[u, v](d(uv) - ud(v) - vd(u)) = 0$, for all $u, v \in U$.

Proof. (i) Since $uv + vu = (u + v)^2 - u^2 - v^2$, we find that $uv + vu \in U$ for all $u, v \in U$. Hence our hypothesis yields the required result.

(ii) Since $uv + vu \in U$, replacing v by $uv + vu$ in (i), we get

$$(2.1) \quad d(u(uv + vu) + (uv + vu)u) = 4u^2d(v) + 6uvd(u) + 2vud(u)$$

On the other hand

$$\begin{aligned} d(u(uv + vu) + (uv + vu)u) &= d(u^2v + vu^2) + 2d(uvu) \\ &= 2u^2d(v) + 4uvd(u) + 2d(uvu) \end{aligned}$$

Combining the above equation with (2.1) we get (ii)

(iii) By linearizing (ii) on u , we get

$$(2.2) \quad \begin{aligned} d((u + w)v(u + w)) &= u^2d(v) + w^2d(v) + (uw + wu)d(v) \\ &\quad + 3uvd(u) + 3wvd(u) + 3uvd(w) \\ &\quad + 3wvd(w) - vud(u) - vud(w) \\ &\quad - vwd(u) - vwd(w) \end{aligned}$$

On the other hand

$$(2.3) \quad \begin{aligned} d((u + w)v(u + w)) &= d(uvu) + d(wvw) + d(uvw + wvu) \\ &= u^2d(v) + 3uvd(u) - vud(u) + w^2d(v) \\ &\quad + 3wvd(w) - vwd(w) + d(uvw + wvu) \end{aligned}$$

Combining (2.2) and (2.3), we get the result.

(iv) Since $uv + vu$ and $uv - vu$ both belong to U , we find that $2uv \in U$ for all $u, v \in U$. Hence, by our hypothesis we find that $d(uv)^2 = 2uvd(uv)$. Replace w

by $2uv$ in (iii), and use the fact that $\text{char } R \neq 2$, to get

$$(2.4) \quad \begin{aligned} d(uv(uv) + (uv)vu) &= (u^2v + uvu)d(v) + 3uvd(uv) \\ &+ 3uv^2d(u) - vud(uv) - vuvd(u) \end{aligned}$$

On the other hand

$$(2.5) \quad \begin{aligned} d((uv)uv + (uv)vu) &= d((uv)^2 + uv^2u) \\ &= 2uvd(uv) + 2u^2vd(v) + 3uv^2d(u) - v^2ud(u) \end{aligned}$$

Combining (2.4) and (2.5), we get

$$(2.6) \quad [u, v]d(uv) = u[u, v]d(v) + v[u, v]d(u), \quad \text{for all } u, v \in U.$$

Replacing $u + v$ for v in (2.6), we have

$$2[u, v]ud(u) + [u, v]d(uv) = 2u[u, v]d(u) + u[u, v]d(v) + v[u, v]d(u).$$

Now application of (2.6) yields the required result.

(v) Linearize (iv) on u , to get

$$\begin{aligned} [u, v]ud(u) + [u, v]vd(v) + [u, v]ud(v) + [u, v]vd(u) \\ = u[u, v]d(u) + u[u, v]d(v) + v[u, v]d(u) + v[u, v]d(v), \quad \text{for all } u, v \in U. \end{aligned}$$

Application of (iv) and (2.6) yield that $[u, v]ud(v) + [u, v]vd(u) = [u, v]d(uv)$ and hence $[u, v]\{d(uv) - ud(v) - vd(u)\} = 0$, for all $u, v \in U$. \square

Lemma 2.3. *Let R be a 2-torsion free ring and let U be a Lie ideal of R such that $u^2 \in U$ for all $u \in U$. If $d : R \rightarrow R$ is an additive mapping satisfying $d(u^2) = 2ud(u)$ for all $u \in U$, then*

- (i) $[u, v]d([u, v]) = 0$, for all $u, v \in U$.
- (ii) $(u^2v - 2uvu + vu^2)d(v) = 0$, for all $u, v \in U$.

Proof. (i) From Lemma 2.2 (i) and (v), we have

$$d(uv + vu) = 2\{ud(v) + vd(u)\} \text{ and } [u, v](d(uv) - ud(v) - vd(u)) = 0$$

respectively. Combining these two results we find that

$$(2.7) \quad [u, v](d(vu) - ud(v) - vd(u)) = 0, \quad \text{for all } u, v \in U.$$

Further, combining of (2.7) and Lemma 2.2 (v) yields that $[u, v]d([u, v]) = 0$.

(ii) For any $v, u \in U$, we have $d([u, v]^2) = 2[u, v]d([u, v])$. Now application of Lemma 2.3 (i), gives that

$$(2.8) \quad d([u, v]^2) = 0, \quad \text{for all } u, v \in U.$$

Since $2uv \in U$, replacing u by $2vu$ in $uv + vu \in U$ and $uv - vu \in U$ and adding the results so obtained we find that $4vuv \in U$ for all $u, v \in U$. Thus in view of Lemma 2.2 (i), we have

$$4d(u(uvu) + (vuv)u) = 8\{ud(vuv) + vuvd(u)\}$$

This implies that $d(u(vuv) + (vuv)u) = 2\{ud(vuv) + vuvd(u)\}$. Now application of (2.8) yields that

$$\begin{aligned} 0 &= d([u, v]^2) \\ &= d(u(vuv) + (vuv)u) - d(uv^2u) - d(vu^2v) \\ &= 2\{ud(vuv) + vuvd(u)\} - u^2d(v^2) - 3uv^2d(u) \\ &\quad + v^2ud(u) - v^2d(u^2) - 3vu^2d(v) + u^2vd(v) \\ &= -3(u^2v - 2uvu + vu^2)d(v) - (uv^2 - 2vuv + v^2u)d(u) \end{aligned}$$

and hence,

$$(2.9) \quad (uv^2 - 2vuv + v^2u)d(u) + 3(u^2v - 2uvu + vu^2)d(v) = 0, \quad \text{for all } u, v \in U.$$

In view of Lemma 2.2 (iv), we have

$$(2.10) \quad (u^2v - 2uvu + vu^2)d(u) = 0, \quad \text{for all } u, v \in U.$$

Replacing u by $u + v$ in (2.10), we find that

$$\{(u^2v - 2uvu + vu^2) - (v^2u - 2vuv + uv^2)\}(d(u) + d(v)) = 0, \quad \text{for all } u, v \in U.$$

Now, using (2.10) in the above expression, we have

$$(2.11) \quad (u^2v - 2uvu + vu^2)d(v) - (v^2u - 2vuv + uv^2)d(u) = 0, \quad \text{for all } u, v \in U.$$

Combining (2.9) and (2.11), and using the fact that R is 2-torsion free, we obtain $(u^2v - 2uvu + vu^2)d(v) = 0$. Thus in view of (2.11), we get the required result. \square

3. MAIN RESULT

The main result of the present paper states as follows:

Theorem. *Let R be a 2-torsion free prime ring and let U be a Lie ideal of R such that $u^2 \in U$. If $d : R \rightarrow R$ is an additive mapping such that $d(u^2) = 2ud(u)$ for all $u \in U$, then $d(uv) = ud(v) + vd(u)$ for all $u, v \in U$.*

Proof. If U is a commutative Lie ideal of R , then by using the same arguments as used in the proof of Lemma 1.3 of [8], $U \subset Z$. Hence using Lemma 2.2 (i), we find that $2d(uv) = 2\{ud(v) + vd(u)\}$. But since $\text{char } R \neq 2$, we find that $d(uv) = ud(v) + vd(u)$ for all $u, v \in U$. Hence onward we shall assume that U is a noncommutative Lie ideal of R - i.e. $U \not\subset Z$. \square

Now, by Lemma 2.2 (iv), we have

$$(3.1) \quad (u^2v - 2uvu + vu^2)d(u) = 0, \quad \text{for all } u, v \in U.$$

Replacing u by $[u_1, w]$ in (3.1), we get

$$([u_1, w]^2v)d([u_1, w]) - 2([u_1, w]v[u_1, w])d([u_1, w]) + (v[u_1, w]^2)d([u_1, w]) = 0,$$

for all $u, v, u_1, w \in U$.

Now, application of Lemma 2.3 (i), yields that $[u_1, w]^2Ud([u_1, w]) = 0$. Hence by Lemma 2.1 either $[u_1, w]^2 = 0$ or $d([u_1, w]) = 0$. If for some $u_1, w \in U$, $d([u_1, w]) = 0$ - i.e. $d(u_1w) = d(wu_1)$, then by using Lemma 2.2 (i) and the fact that

char $R \neq 2$, we get $d(u_1w) = u_1d(w) + wd(u_1)$. On the other hand let $[u_1, w]^2 = 0$, for some $u_1, w \in U$. By Lemma 2.3 (ii), we get

$$(3.2) \quad (u^2v - 2uvu + vu^2)d(v) = 0, \quad \text{for all } u, v \in U.$$

Replacing v by $[u_1, w]$ in (3.2), we get

$$(u^2[u_1, w])d([u_1, w]) - 2(u[u_1, w]u)d([u_1, w]) + ([u_1, w]u^2)d([u_1, w]) = 0,$$

for all $u \in U$.

Again apply Lemma 2.3 (i), to get

$$(3.3) \quad ([u_1, w]u^2)d([u_1, w]) - 2(u[u_1, w]u)d([u_1, w]) = 0, \quad \text{for all } u \in U.$$

Linearizing (3.3) on u and using (3.2), we have

$$(3.4) \quad ([u_1, w]uv)d([u_1, w]) + ([u_1, w]vu)d([u_1, w]) - 2\{(u[u_1, w]v) + (v[u_1, w]u)\}d([u_1, w]) = 0, \quad \text{for all } u, v \in U.$$

Replace u by $2uv_1$ in (3.4) and use the fact that R is 2-torsion free, to get

$$([u_1, w]uv_1v)d([u_1, w]) + ([u_1, w]vuv_1)d([u_1, w]) - 2\{(uv_1[u_1, w]v) + (v[u_1, w]uv_1)\}d([u_1, w]) = 0, \quad \text{for all } u, v, v_1 \in U.$$

Further, replacing v_1 by $[u_1, w]$ in the above expression and applying Lemma 2.3 (i) together with the fact that $[u_1, w]^2 = 0$, we find that $([u_1, w]u[u_1, w])v d([u_1, w]) = 0$ - i.e. $([u_1, w]u[u_1, w])Ud([u_1, w]) = 0$, for all $u \in U$. Thus by Lemma 2.1 either $d([u_1, w]) = 0$ or $[u_1, w]u[u_1, w] = 0$. If $d([u_1, w]) = 0$, then using the similar arguments as above we get the required result. On the other hand if $[u_1, w]u[u_1, w] = 0$ for all $u \in U$, then again by Lemma 2.1 we have $[u_1, w] = 0$. Further, application of Lemma 2.2 (i) yields that $2d(u_1w) = 2\{u_1d(w) + wd(u_1)\}$ and hence $d(u_1w) = u_1d(w) + wd(u_1)$. Hence in both the cases we find that $d(uv) = ud(v) + vd(u)$, for all $u, v \in U$. This completes the proof of the above theorem.

Corollary. *Let R be a 2-torsion free prime ring and $d : R \rightarrow R$ be a Jordan left derivation. Then d is a left derivation.*

Acknowledgement. The authors are thankful to Prof. Murtaza A. Quadri for encouragement and fruitful discussion.

REFERENCES

[1] Awtar, R., *Lie ideals and Jordan derivations of prime rings*, Proc. Amer. Math. Soc. **90** (1984), 9-14.
 [2] Bergen, J., Herstein, I. N. and Ker, J. W., *Lie ideals and derivations of prime rings*, J. Algebra **71** (1981), 259-267.
 [3] Bresar, M., *Jordan derivations on semiprime rings*, Proc. Amer. Math. Soc. **104** (1988), 1003-1006.
 [4] Bresar, M. and Vukman, J., *Jordan derivations of prime rings*, Bull. Aust. Math. Soc. **37** (1988), 321-322.

- [5] Bresar, M. and Vukman, J., *On left derivations and related mappings*, Proc. Amer. Math. Soc. **110** (1990), 7–16.
- [6] Deng, Q., *On Jordan left derivations*, Math. J. Okayama Univ. **34** (1992), 145–147.
- [7] Herstein, I.N., *Jordan derivations of prime rings*, Proc. Amer. Math. Soc. **8** (1957), 1104–1110.
- [8] Herstein, I.N., *Topics in ring theory*, Univ. of Chicago Press, Chicago 1969.
- [9] Kill-Wong Jun and Byung-Do Kim, *A note on Jordan left derivations*, Bull. Korean Math. Soc. **33** (1996) No. 2, 221–228.
- [10] Posner, E.C., *Derivations in prime rings*, Proc. Amer. Math. Soc. **8** (1957), 1093–1100.
- [11] Vukman, J., *Jordan left derivations on semiprime rings*, Math. J. Okayama Univ. (to appear).

M. ASHRAF, DEPARTMENT OF MATHEMATICS
FACULTY OF SCIENCE, KING ABDUL AZIZ UNIVERSITY
P.O. BOX. 9028, JEDDAH 21413, SAUDI-ARABIA

NADEEM-UR-REHMAN, DEPARTMENT OF MATHEMATICS
ALIGARH MUSLIM UNIVERSITY
ALIGARH 202002, INDIA